


CLAIMS


1. Process for preparing a pharmaceutically active 5 compound, zolmitriptan, or a pharmaceutically acceptable salt thereof, **characterised** in that it comprises the following stages:

a) Preparation of the diazononium salt from the 10 aniline hydrochloride of formula (II)



(II)

15 followed by reduction and acidification to give the hydrazine of formula (III):




(III)

b) *In situ* reaction of the hydrazine hydrochloride of formula (III) with α -keto- δ -valerolactone, to give the hydrazone of formula (IV):


ART 34 AMDT

21




(IV)

c) Fischer indole synthesis of the hydrazone of formula (IV), to give the pyranoindolone of formula (V):



(V)

10 d) Transesterification of the pyranoindolone of formula (V), to provide the compound of formula (VI):



(VI)


15

in which R represents a straight or branched C1-C4 alkyl chain;

e) Conversion of the hydroxyl group of the compound of formula (VI) into dimethylamino, to give the indolecarboxylate of formula (VII):

ART 34 AMDT

22




(VII)

5

in which R represents a straight or branched C1-C4 alkyl chain;

f) Saponification of the 2-carboalkoxy group of the compound of formula (VII), to give the indolecarboxylic acid of formula (VIII):



(VIII)

15

g) Decarboxylation of the indolecarboxylic acid of formula (VIII), to give zolmitriptan and,

eventually, the preparation of a pharmaceutically acceptable salt thereof.

2. Process as claimed in Claim 1, characterised in that said stage c) is carried out in a solution of dry hydrogen chloride in acetic acid.

BEST AVAILABLE COPY

ART 34 AMDT

23

3. Process as claimed in Claim 1, characterised in that said stages c) and d) are carried out in a one pot reaction.

5

4. Process as claimed in Claim 1 and Claim 3, characterised in that said stages c) and d) are carried out in a solution of dry hydrogen chloride in a straight or branched C1-C4 alcohol chain.

10


5. Process as claimed in Claim 1, characterised in that said stage e) is carried out in two steps:

e-i) replacement of the hydroxyl group of the compound of formula (VI) by a leaving group X; and

15 e-ii) subsequent substitution reaction of the leaving group X with dimethylamine to provide the compound of formula (VII).

6. Process as claimed in Claim 5, characterised in that said leaving group X is chosen between an atom of halogen, a mesylate group or a tosylate group.

7. Synthesis intermediate of formula (IV):




25

(IV)

ART 34 AMDT


24

8. Synthesis intermediate of formula (V):



5


9. Synthesis intermediate of formula (VI):



10

where R represents a straight or branched C1-C4 alkyl
15 chain.

10. Synthesis intermediate of formula (VIII):



20